

MOMENTS OF THE BLOCK OPERATORS IN THE GROUP VON NEUMANN ALGEBRAS

ILWOO CHO

ABSTRACT. In this paper, we will consider the moments of the block operators of the given group von Neumann algebra $L(G)$, where the given group G is a finitely presented discrete group $\langle X : R \rangle$, where X is the generator set of G and R is the relation on the set X , as the set of relators. Define the canonical trace tr on $L(G)$ and the W^* -probability space $(L(G), tr)$ which is our free probabilistic object of this paper. Define the block operators T_x by $T_x = x + x^{-1}$ of $L(G)$, where $x \in X$. In this paper, we will compute the moments and the moment series of T_x , for $x \in X$. By the computation, we can get that if x_1 and x_2 are generators of presented groups $\langle X_1 : R_1 \rangle$ and $\langle X_2 : R_2 \rangle$, respectively, and (i) if there is $n \in \mathbb{N}$ such that $x_1^n \in R_1$ and $x_2^n \in R_2$, or (ii) if there is no $n_1, n_2 \in \mathbb{N}$ such that $x_1^{n_1} \in R_1$ and $x_2^{n_2} \in R_2$, then the block operators $x_1 + x_1^{-1}$ in $L(G_1)$ and $x_2 + x_2^{-1}$ in $L(G_2)$ are identically distributed.

The group von Neumann algebras are studied recently by various authors. Group von Neumann algebras are interesting objects in Operator Algebra and Free Probability. In this paper, we will consider the moments of certain operators in group von Neumann algebras, where the group is presented by a finite generator set and a finite relation. We will take a presented group $\langle X : R \rangle$, where X is the generator set and R is the relation on the group, as the nonempty set of relators. For instance, the symmetric group S_3 can be presented by its generator set X_{S_3} and its relation R_{S_3} , where

$$X_{S_3} = \{a, b\}$$

and

$$R_{S_3} = \{a^2, b^3, (ab)^2\}.$$

Let $H = \langle X : R \rangle$ be a presented group and let $L(H)$ be the group von Neumann algebra generated by H . i.e.,

$$L(H) = \overline{\lambda(H)}^w = \overline{\mathbb{C}[H]}^w,$$

where λ is the left (unitary) representation. If $a \in L(H)$ is an operator, then it has the Fourier expansion,

$$a = \sum_{h \in H} a_h h, \text{ for } a_h \in \mathbb{C} \text{ and } h \in H.$$

Key words and phrases. Finitely Presented Groups, Group von Neumann Algebras, W^* -Probability Spaces, Moments of Random Variables.

In fact, the group element h in the previous expansion are understood as unitary operators λ_h on the Hilbert space $l^2(H)$. Recall that

$$x^* = \sum_{h \in H} \overline{a_h} h^{-1},$$

where $h^{-1} = \lambda_{h^{-1}} = \lambda_h^* = h^*$, on $l^2(H)$, for all $h \in H$. Define the canonical trace tr on $L(H)$ by

$$tr(x) = tr\left(\sum_{h \in H} a_h h\right) = a_{e_H}, \text{ for all } x \in L(H)$$

where e_H is the identity of H . Then we can have the W^* -probability space $(L(H), tr)$. The main purpose of this paper is to compute the moments of the block operators of $L(H)$. In order to do that we observed the free monoid $X^* = (X \cup X^{-1})'$ of the group H and the corresponding combinatorial forms in X^* of the group elements in G . (If Y is an arbitrary set, then Y' is the set of all free words in Y , which is called the free monoid of Y . The elements in Y' are called the combinatorial forms of $\langle Y \rangle$, where $\langle Y \rangle$ is the group generated by Y .)

Let $G = \langle X : R \rangle$ be a presented group with its generator set $X = \{x_1, \dots, x_N\}$ and its relation $R = \{r_1, \dots, r_M\}$. In Chapter 1, we will consider the free monoid X^* of the group G defined by

$$X^* \stackrel{def}{=} \bigcup_{n=0}^{\infty} \left\{ x_{j_1}^{p_1} \dots x_{j_n}^{p_n} : \begin{array}{l} (j_1, \dots, j_n) \in \{1, \dots, N\}^n \\ (p_1, \dots, p_n) \in \{1, -1\}^n \end{array} \right\},$$

which is the set of all free words of the generator set X and X^{-1} . When $n = 0$, the corresponding word is the empty word $\emptyset$. There exists a monoid homomorphism π from X^* onto the given group G . Notice that, for any $g \in G$, there is a subset $\pi^{-1}(g)$ in X^* . The elements w_g in $\pi^{-1}(g)$ are called the combinatorial forms of $g \in G$. We will use the word problem on X^* , by computing the moments of the block operators.

In Chapter 2, we will compute the moments of the block operators $T_x = x + x^{-1}$, for $x \in X$, in the W^* -probability space $(L(G), tr)$. In particular, we have that ;

(1) Suppose that there is no relator $r_t \in R$ such that $x^k = r_t$, for all $k \in \mathbb{N}$. Then

$$tr(T_x^m) = \left[\begin{array}{c} m \\ \frac{m}{2} \end{array} \right], \text{ for all } m \in \mathbb{N},$$

$$\text{where } \left[\begin{array}{c} m \\ \frac{m}{2} \end{array} \right] \stackrel{def}{=} \begin{cases} \left(\begin{array}{c} m \\ \frac{m}{2} \end{array} \right) = \frac{m!}{(\frac{m}{2}!)^2} & \text{if } m \text{ is even} \\ 0 & \text{if } m \text{ is odd.} \end{cases}$$

(2) Suppose that there exist $r_t \in R$ and $n_x \in \mathbb{N}$ such that $r_t = x^{n_x}$. Then

$$tr(T_x^m) = \begin{cases} \left[\begin{array}{c} m \\ \frac{m}{2} \end{array} \right] & \text{if } m < n_j \\ \left(2^{k_1} - \left[\begin{array}{c} k_1 \\ \frac{k_1}{2} \end{array} \right] \right) + \left[\begin{array}{c} m \\ \frac{m}{2} \end{array} \right] & \text{if } m = k_1 n_j + k_2, \end{cases}$$

where $k_1 \in \mathbb{N}$ and $k_2 \in \mathbb{N} \cup \{0\}$ such that $0 \leq k_2 < n_x$.

Notice that if G_1 and G_2 are finitely presented groups (not necessarily distinct) and x_1 and x_2 are generators of G_1 and G_2 , respectively, and if there exists $n \in \mathbb{N}$ such that x_1^n and x_2^n are relators of G_1 and G_2 , respectively, then the block operators $x_1 + x_1^{-1}$ in $L(G_1)$ and $x_2 + x_2^{-1}$ in $L(G_2)$ are identically distributed. Also, if x_1 and x_2 have no relators x_1^n and x_2^k , for $n, k \in \mathbb{N}$, then they are also identically distributed.

1. PRELIMINARIES

In this paper, we will compute the moments of certain operators on a group von Neumann algebra with its canonical faithful normal trace. We will restrict our interests to finitely presented group von Neumann algebras. Let $G = \langle X : R \rangle$ be a presented group, where X is the finite generator set of the group G and R is the relation on G , as the set of all relators. Denote the corresponding group von Neumann algebra by $L(G)$. Then each operator $a \in L(G)$ has its Fourier expansion

$$(1.1) \quad a = \sum_{g \in G} a_g g, \text{ for } a_g \in \mathbb{C}.$$

Note that we can regard g in (1.3) as λ_g , for all $g \in G$, where λ is the left regular representation. Remark that $g^* = g^{-1}$, for all $g \in G$, in $L(G)$, and hence each operator $g \in L(G)$ is unitary. For the group von Neumann algebra $L(G)$, we can define the canonical trace tr by

$$(1.2) \quad tr(a) \stackrel{def}{=} tr \left(\sum_{g \in G} a_g g \right) = a_{e_G},$$

for all $a \in L(G)$, where e_G is the identity of the group G .

Definition 1.1. Let $G = \langle X : R \rangle$ be the presented group and $L(G)$, the corresponding group von Neumann algebra. The algebraic pair $(L(G), tr)$ is called the presented group W^* -probability space, where tr is the canonical trace given in (1.2).

The operators in $(L(G), tr)$ are called the random variables. Let $a \in L(G)$. Then the n -th moments of a is defined by

$$tr(a^n), \quad \text{for all } n \in \mathbb{N}.$$

Now, we have our free probabilistic objects of this paper.

2. MOMENTS OF THE BLOCK OPERATORS IN GROUP VON NEUMANN ALGEBRAS

Throughout this chapter, let $G = \langle X : R \rangle$ be the fixed presented graph with its generator set X and its relation R

$$X = \{x_1, \dots, x_N\}$$

and

$$R = \{r_1, \dots, r_M\},$$

where $M, N \in \mathbb{N}$. In this chapter, we will compute the moments of block operators $T_x = x + x^{-1}$, for $x, x^{-1} \in X$, for $j = 1, \dots, N$.

Definition 2.1. Let $G = \langle X : R \rangle$ be a presented group with its generator set $X = \{x_1, \dots, x_N\}$ and its relation $R = \{r_1, \dots, r_M\}$, where $M, N \in \mathbb{N}$ and $r_1, \dots, r_M$ are relators, as elements in X^* . The operators T_j are block operators in the group von Neumann algebra $L(G)$, if

$$T_j = x_j + x_j^{-1}, \quad \text{for all } j = 1, \dots, N.$$

2.1. Moments of Block Operators.

Let Y be an arbitrary set. Then we can define a set Y' , consisting of all free words in Y . This set Y' is called the free set of Y . Let $G = \langle X : R \rangle$ be the finitely presented group with

$$X = \{x_1, \dots, x_N\} \quad \text{and} \quad R = \{r_1, \dots, r_M\}.$$

Define the set X^* be the free monoid $(X \cup X^{-1})'$ of the set $X \cup X^{-1}$. Notice that, there exists the surjective (monoid) homomorphism $\pi : X^* \rightarrow G$ and, for any group element g in G , there exist words w_g in X^* satisfying that $\pi(w_g) = g$ in G . It is easy to see that a corresponding word w_g of g is not uniquely determined. We

say that such words $w_g \in \pi^{-1}(g)$ of $g \in G$ are combinatorial forms of g . Let $w_g = x_{j_1}^{p_1} \dots x_{j_n}^{p_n} \in X^*$ be a combinatorial form of $g \in G$, where

$$(j_1, \dots, j_n) \in \{1, \dots, N\}^n \text{ and } (p_1, \dots, p_n) \in \{\pm 1\}^n.$$

For convenience, we denote the word $x_{j_n}^{-p_n} \dots x_{j_1}^{-p_1}$ by w_g^{-1} . Notice that $w_g^{-1} \in \pi^{-1}(g^{-1})$ in X^* .

Now, fix the generators $x_j \in X$ and the block operator $T_j = x_j + x_j^{-1}$. Consider the n -th moments $\text{tr}(T_j^m)$ of T_j , for all $m \in \mathbb{N}$. It is easy to see that

$$(2.1) \quad T_j^m = \sum_{(p_1, \dots, p_m) \in \{1, -1\}^m} (x_j^{p_1} \dots x_j^{p_m}) = \sum_{(p_1, \dots, p_m) \in \{1, -1\}^m} x_j^{\sum_{k=1}^m p_k},$$

for all $m \in \mathbb{N}$. Notice that each word $x_j^{\sum_{k=1}^m p_k} = x_j^{p_1} \dots x_j^{p_m}$ is regarded as an element in the free monoid X^* . i.e., without loss of generality, we can consider the summands of T_j^m as elements in the free monoid X^* .

Lemma 2.1. *Let $T_j = x_j + x_j^{-1}$ be a block operator of the generator $x_j \in X$. If there is no $n_j \in \mathbb{N}$ such that $x_j^{n_j} \in R$, then*

$$\text{tr}(T_j^m) = \begin{cases} \binom{m}{\frac{m}{2}} & \text{if } m \in 2\mathbb{N} \\ 0 & \text{if } m \in 2\mathbb{N} - 1, \end{cases}$$

$$\text{where } \binom{n}{k} = \frac{n!}{k!(n-k)!}, \text{ for } n, k \in \mathbb{N}.$$

Proof. Assume that there is no relator $r_t \in R$ and $n_j \in \mathbb{N}$ such that $r_t = x_j^{n_j}$ in $R \subset X^*$. Then, by (2.1), we have that

$$\begin{aligned} \text{tr}(T_j^m) &= \text{tr} \left(\sum_{(p_1, \dots, p_m) \in \{1, -1\}^m} x_j^{\sum_{k=1}^m p_k} \right) \\ &= \text{tr} \left(\sum_{(p_1, \dots, p_m), \sum_{j=1}^m p_j = 0} x_j^{\sum_{k=1}^m p_k} \right) \\ &= \text{tr} \left(\sum_{(p_1, \dots, p_m), \sum_{k=1}^m p_j = 0} e_G \right) \\ &= |\{(p_1, \dots, p_m) \in \{1, -1\}^n : \sum_{k=1}^n p_j = 0\}| \end{aligned}$$

$$= \begin{pmatrix} m \\ \frac{m}{2} \end{pmatrix}.$$

The last equality holds because, to make $\sum_{k=1}^m p_k = 0$, the same number of $+1$'s and -1 's should be appeared in the sequence $(p_1, \dots, p_m)$. It is easy to see that if m is odd, then the set

$$\{(p_1, \dots, p_m) \in \{1, -1\}^m : \sum_{k=1}^m p_k = 0\}$$

is empty. Therefore, all odd moments of T_j vanish. ■

Now, we assume that there is a relator $r_t \in R$ such that $r_t = x_j^{n_j}$, for $n_j \in \mathbb{N}$, as a free word in X^* . Clearly, the relator r_t is a combinatorial form of e_G (i.e., $\pi(r_t) = e_G$) in the group G and the length $|r_t|$ of r_t in X^* is n_j . Also, notice that if $r_t \in R$, then the words $wr_t w^{-1}$ and $wr_t^{-1} w^{-1}$ are also combinatorial forms of e_G , for all words w and w in X^* . By (2.1), we have that

$$T_j^m = \sum_{(p_1, \dots, p_m) \in \{1, -1\}^m} x_j^{\sum_{k=1}^m p_k}.$$

If $m = n_j$, then

$$\begin{aligned} T_j^{n_j} &= \left(x_j^{n_j} + x_j^{-n_j} \right) + \sum_{(p_1, \dots, p_{n_j}) \in \{1, -1\}^{n_j}, (p_1, \dots, p_{n_j}) \neq (\pm 1, \dots, \pm 1)} x_j^{\sum_{k=1}^{n_j} p_k} \\ (2.2) \quad &= 2e_G + \sum_{(p_1, \dots, p_{n_j}) \in \{1, -1\}^{n_j}, (p_1, \dots, p_{n_j}) \neq (\pm 1, \dots, \pm 1)} x_j^{\sum_{k=1}^{n_j} p_k}. \end{aligned}$$

By the above formula (2.2), we can get the following lemma ;

Lemma 2.2. *Let $x_j \in X$ and assume that there exists $t \in \{1, \dots, M\}$ such that $r_t = x_j^{n_j}$, for $n_j \in \mathbb{N}$. Then*

$$\begin{aligned} \text{tr}(T_j^m) &= \begin{cases} \begin{pmatrix} m \\ \frac{m}{2} \end{pmatrix} & \text{for all even } m < n_j \\ 0 & \text{for all odd } m < n_j \end{cases} \\ \text{and} \\ \text{tr}(T_j^{n_j}) &= \begin{cases} 2 & \text{if } n_j \in 2\mathbb{N} - 1 \\ 2 + \begin{pmatrix} n_j \\ \frac{n_j}{2} \end{pmatrix} & \text{if } n_j \in 2\mathbb{N}. \end{cases} \end{aligned}$$

Proof. The first formula is trivial, by the previous lemma.

Suppose that n_j is an odd number in $\mathbb{N}$. Then, by (2.2), we have that

$$T_j^{n_j} = 2e_G + \sum_{(p_1, \dots, p_{n_j}) \in \{1, -1\}^{n_j}, (p_1, \dots, p_{n_j}) \neq (\pm 1, \dots, \pm 1)} x_j^{\sum_{k=1}^{n_j} p_k}.$$

Since n_j is an odd number, we cannot find the sequence $(p_1, \dots, p_{n_j})$ in $\{\pm 1\}^{n_j}$ satisfying that $\sum_{k=1}^{n_j} p_k = 0$. So, we cannot find the e_G -terms in the summand

$$\sum_{(p_1, \dots, p_{n_j}) \in \{1, -1\}^{n_j}, (p_1, \dots, p_{n_j}) \neq (\pm 1, \dots, \pm 1)} x_j^{\sum_{k=1}^{n_j} p_k}$$

of $T_j^{n_j}$. Thus if n_j is an odd number, then $\text{tr}(T_j^{n_j}) = 2$.

Now, assume that n_j is an even number in $\mathbb{N}$. Then, again by (2.2), we have that

$$\begin{aligned} \text{tr}(T_j^{n_j}) &= 2 + \sum_{(p_1, \dots, p_{n_j}) \in \{1, -1\}^{n_j}, (p_1, \dots, p_{n_j}) \neq (\pm 1, \dots, \pm 1)} \text{tr}\left(x_j^{\sum_{k=1}^{n_j} p_k}\right) \\ &= 2 + |\{(p_1, \dots, p_{n_j}) \in \{\pm 1\}^{n_j} : \sum_{k=1}^{n_j} p_k = 0\}| \\ &= 2 + \binom{n_j}{\frac{n_j}{2}}. \end{aligned}$$

■

Now, suppose that $m > n_j$. There are two cases ;

- (i) $m = k_1 n_j + k_2$, where $1 \leq k_2 < n_j$ or
- (ii) $m = k n_j$, for some $k \in \mathbb{N}$.

Lemma 2.3. *Let $x_j \in X$ and assume that there exists $t \in \{1, \dots, M\}$ such that $r_t = x_j^{n_j}$, for $n_j \in \mathbb{N}$. Then*

(1) *If $m = k n_j$, for $k \in \mathbb{N}$, then*

$$\text{tr}(T_j^m) = \left(2^k - \left\lfloor \frac{k}{2} \right\rfloor\right) + \left\lfloor \frac{k n_j}{2} \right\rfloor.$$

(2) *If $m = k_1 n_j + k_2$, for $k_1, k_2 \in \mathbb{N}$ and if $1 \leq k_2 < n_j$, then*

$$\text{tr}(T_j^m) = 2^{k_1} + \left\lfloor \frac{m}{2} \right\rfloor - \left\lfloor \frac{k_1}{2} \right\rfloor,$$

where

$$\begin{bmatrix} t \\ \frac{t}{2} \end{bmatrix} \stackrel{def}{=} \begin{cases} \begin{pmatrix} t \\ \frac{t}{2} \end{pmatrix} & \text{if } t \text{ is even} \\ 0 & \text{if } t \text{ is odd.} \end{cases}$$

Proof. (1) Let $(p_1, \dots, p_m) \in \{\pm 1\}^m$, where $m = kn_j$ is sufficiently big number in $\mathbb{N}$, where $k, n_j \in \mathbb{N}$. Define subsequences

$$\mathbf{i}_+ = \left(\underbrace{1, \dots, 1}_{n_j\text{-times}} \right) \quad \text{and} \quad \mathbf{i}_- = \left(\underbrace{-1, \dots, -1}_{n_j\text{-times}} \right).$$

Since $m = kn_j$, there exists a sequence $P = (p_1, \dots, p_m)$ such that

$$P = (\mathbf{i}_{i_1}, \dots, \mathbf{i}_{i_k}), \text{ for } i_1, \dots, i_k \in \{+, -\}.$$

We define the set W_j , consisting of such sequences. i.e.,

$$(2.3) \quad W_j \stackrel{def}{=} \{(\mathbf{i}_{i_1}, \dots, \mathbf{i}_{i_k}) : i_1, \dots, i_k \in \{+, -\}\}.$$

Note that, for $\mathbf{i}_+$, we have $x_j^{n_j} = r_{t_j}$ and, for $\mathbf{i}_-$, we have $x_j^{-n_j} = r_{t_j}^{-1}$ in X^* . Thus, we can get that

$$(2.4) \quad |W_j| = 2^k \quad \text{and}$$

$$(2.5) \quad \sum_{k=1}^m p_k = pn_j, \text{ where } p = 1 \text{ or } \dots \text{ or } k,$$

for all $(p_1, \dots, p_m)$ in W_j . We will define a subset W'_j of the set W_j (if exists) by

$$(2.6) \quad W'_j = \left\{ (\mathbf{i}_{i_1}, \dots, \mathbf{i}_{i_k}) : \begin{array}{l} i_{j_1} = \dots = i_{j_{\frac{k}{2}}} = +, \\ i_{s_1} = \dots = i_{s_{\frac{k}{2}}} = -, \\ \{i_{j_1}, \dots, i_{j_{\frac{k}{2}}}\} \cup \{i_{s_1}, \dots, i_{s_{\frac{k}{2}}}\} = \{i_1, \dots, i_k\} \end{array} \right\}$$

Then $W'_j \subseteq W_j$ and, since there are same numbers of +’s and -’s, we can have

$$(2.7) \quad |W'_j| = \binom{k}{\frac{k}{2}},$$

if the nonempty subset W'_j exists in W_j . (It is easily see that if k is even, then W'_j exists in W_j . And if k is odd, then W'_j is empty.) Define the subset S_0 of sequences in $\{\pm 1\}^m$ by

$$S_0 = \{(p_1, \dots, p_m) \in \{\pm 1\}^m : \sum_{k=1}^m p_k = 0\}.$$

Then, by (2.6), $W'_j \subset S_0$. In fact,

$$(2.8) \quad S_0 \cup W'_j = S_0 \text{ and } (W_j \setminus W'_j) \cap S_0 = \emptyset.$$

Assume that k is even. Then

$$\begin{aligned} \text{tr}(T_j^m) &= \text{tr} \left(\sum_{(p_1, \dots, p_m) \in \{\pm 1\}^m} x_j^{\sum_{k=1}^m p_k} \right) \\ &= \text{tr} \left(\sum_{(p_1, \dots, p_m) \in W_j \setminus W'_j} x_j^{\sum_{k=1}^m p_k} \right) \\ &\quad + \text{tr} \left(\sum_{(p_1, \dots, p_m) \in \{\pm 1\}^m, (p_1, \dots, p_m) \in S_0} x_j^{\sum_{k=1}^m p_k} \right) \\ &= |W_j \setminus W'_j| + \text{tr} \left(\sum_{(p_1, \dots, p_m) \in \{\pm 1\}^m, (p_1, \dots, p_m) \in S_0} e_G \right) \end{aligned}$$

where W_j is defined in (2.3), by (2.8)

$$\begin{aligned} &= \left(2^k - \binom{k}{\frac{k}{2}} \right) + |S_0| \\ (2.9) \quad &= \begin{cases} \left(2^k - \binom{k}{\frac{k}{2}} \right) + \binom{m}{\frac{m}{2}} & \text{if } m \text{ is even} \\ 2^k - \binom{k}{\frac{k}{2}} & \text{if } m \text{ is odd} \end{cases}, \end{aligned}$$

where $k \in 2\mathbb{N}$ and $m \in \mathbb{N}$. Now, let's suppose that k is an odd number greater than 1 in $\mathbb{N}$. Then we can have that

$$(2.10) \quad W'_j = \emptyset \quad \text{and} \quad W_j \cap S_0 = \emptyset.$$

Therefore, by (2.10), we have that

$$\text{tr}(T_j^m) = \text{tr} \left(\sum_{(p_1, \dots, p_m) \in W_j} x_j^{\sum_{k=1}^m p_k} \right)$$

$$\begin{aligned}
& +tr \left(\sum_{(p_1, \dots, p_m) \in \{\pm 1\}^m, (p_1, \dots, p_m) \in S_0} x_j^{\sum_{k=1}^m p_k} \right) \\
& = |W_j| + tr \left(\sum_{(p_1, \dots, p_m) \in \{\pm 1\}^m, (p_1, \dots, p_m) \in S_0} e_G \right) \\
& = |W_j| + |S_0| \\
(2.11) \quad & = \begin{cases} 2^k + \binom{m}{\frac{m}{2}} & \text{if } m \text{ is even} \\ 2^k & \text{if } m \text{ is odd.} \end{cases}
\end{aligned}$$

Now, define a new notation

$$\left[\begin{array}{c} t \\ \frac{t}{2} \end{array} \right] \stackrel{def}{=} \begin{cases} \binom{t}{\frac{t}{2}} & \text{if } t \in 2\mathbb{N} \\ 0 & \text{if } t \in 2\mathbb{N} - 1. \end{cases}$$

Then the formuli (2.9) and (2.11) can be shortened by

$$tr \left(T_j^{kn_j} \right) = \left(2^k - \left[\frac{k}{2} \right] \right) + \left[\frac{kn_j}{2} \right].$$

(2) Let's assume that $m = k_1 n_j + k_2$, where $n_j \nmid k_2$ and $1 \leq k_2 < n_j$. By (2.1), we have that

$$T_j^m = \sum_{(p_1, \dots, p_m) \in \{\pm 1\}^m} x^{\sum_{k=1}^m p_k}.$$

Let's regard the summands $x^{\sum_{k=1}^m p_k}$ as free words in the free monoid X^* . Then there is a set

$$S_0 = \{(p_1, \dots, p_m) \in \{\pm 1\}^m : \sum_{k=1}^m p_k = 0\}$$

with its cardinality $|S_0| = \left[\frac{m}{2} \right]$. Now, define a set of free words W^j by

$$W^j = \left\{ ((p_{t_1}, \dots, p_{t_{k_2}}) \rightsquigarrow (\mathbf{i}_{i_1}, \dots, \mathbf{i}_{i_{k_1}})) : \begin{array}{l} p_{t_k} \in \{\pm 1\}, \\ i_k \in \{+, -\}, \end{array} \right\},$$

where $\mathbf{i}_+$ and $\mathbf{i}_-$ are defined in (1) and $\rightsquigarrow$ means the insertion. i.e., the sequence

$$((p_{t_1}, \dots, p_{t_{k_2}}) \rightsquigarrow (\mathbf{i}_{i_1}, \dots, \mathbf{i}_{i_{k_1}}))$$

is the free word in X^* with its length $m = k_1 n_j + k_2$. For example,

$$((p_1) \rightsquigarrow (\mathbf{i}_+, \mathbf{i}_+))$$

is

$$(\mathbf{i}_+, p_1, \mathbf{i}_+) \text{ or } (p_1, \mathbf{i}_+, \mathbf{i}_+) \text{ or } (\mathbf{i}_+, \mathbf{i}_+, p_1).$$

where $m = 2n_j + 1$. Define the subset $W_{k_2}^j$ of W^j (if exists) by

$$W_{k_2}^j = \{((p_{t_1}, \dots, p_{t_{k_2}}) \rightsquigarrow (\mathbf{i}_{i_1}, \dots, \mathbf{i}_{i_{k_1}})) : \sum_{i=1}^{k_2} p_{t_i} = 0\}.$$

Let's assume that $W_{k_2}^j$ exists. Then we can define the subset $W_{k_2}^j(S_0)$ of $W_{k_2}^j$ by

$$W_{k_2}^j(S_0) = \left\{ ((p_{t_1}, \dots, p_{t_{k_2}}) \rightsquigarrow (\mathbf{i}_{i_1}, \dots, \mathbf{i}_{i_{k_1}})) : \begin{array}{l} \sum_{i=1}^{k_2} p_{t_i} = 0 \\ (\mathbf{i}_{i_1}, \dots, \mathbf{i}_{i_{k_1}}) \in W_j' \end{array} \right\},$$

where W_j' is defined in (1). It is easy to see that if both k_1 and k_2 are even, then $W_{k_2}^j$ exists in $W_{k_2}^j$. Then

$$(2.12) \quad |W_{k_2}^j| = 2^{k_1} + \left\lceil \frac{k_2}{2} \right\rceil,$$

$$(2.13) \quad |W_{k_2}^j(S_0)| = \left\lceil \frac{k_1}{2} \right\rceil + \left\lceil \frac{k_2}{2} \right\rceil,$$

$$(2.14) \quad W_{k_2}^j(S_0) \cup S_0 = S_0,$$

$$(2.15) \quad |S_0 \setminus W_{k_2}^j(S_0)| = \left\lceil \frac{m}{2} \right\rceil - \left(\left\lceil \frac{k_1}{2} \right\rceil + \left\lceil \frac{k_2}{2} \right\rceil \right)$$

By (2.12) and (2.15), we can compute that ;

$$\begin{aligned} tr(T_j^m) &= tr \left(\sum_{(p_1, \dots, p_m) \in \{\pm 1\}^m} x^{\sum_{k=1}^m p_k} \right) \\ &= tr \left(\sum_{(p_1, \dots, p_n) \in W_{k_2}^j} x^{\sum_{k=1}^m p_k} \right) \\ &\quad + tr \left(\sum_{(p_1, \dots, p_m) \in S_0 \setminus W_{k_2}^j(S_0)} x^{\sum_{k=1}^m p_k} \right) \\ &= |W_{k_2}^j| + |S_0 \setminus W_{k_2}^j(S_0)| \end{aligned}$$

$$\begin{aligned}
&= \left(2^{k_1} + \left[\begin{array}{c} k_2 \\ \frac{k_2}{2} \end{array} \right] \right) + \left(\left[\begin{array}{c} m \\ \frac{m}{2} \end{array} \right] - \left(\left[\begin{array}{c} k_1 \\ \frac{k_1}{2} \end{array} \right] + \left[\begin{array}{c} k_2 \\ \frac{k_2}{2} \end{array} \right] \right) \right) \\
&= 2^{k_1} + \left[\begin{array}{c} m \\ \frac{m}{2} \end{array} \right] - \left[\begin{array}{c} k_1 \\ \frac{k_1}{2} \end{array} \right].
\end{aligned}$$

■

Remark that, by the previous lemma, we have that if $k = 1$, then,

$$\text{tr}(T_j^{n_j}) = \left(2^1 - \left[\begin{array}{c} 1 \\ \frac{1}{2} \end{array} \right] \right) + \left[\begin{array}{c} n_j \\ \frac{n_j}{2} \end{array} \right] = 2 + \left[\begin{array}{c} n_j \\ \frac{n_j}{2} \end{array} \right].$$

By the previous lemmas, we can get the following theorem ;

Theorem 2.4. *Let $G = \langle X : R \rangle$ be a finitely presented group with its generator set $X = \{x_1, \dots, x_N\}$ and its relation $R = \{r_1, \dots, r_M\}$. Fix a generator $x_j \in X$ satisfying that $r_t = x_j^{n_j}$, for some $n_j \in \mathbb{N} \setminus \{1\}$, where $r_t \in R$. Then*

$$(1) \text{ if } m < n_j, \text{ then } \text{tr}(T_j^m) = \left[\begin{array}{c} m \\ \frac{m}{2} \end{array} \right],$$

$$(2) \text{ if } m = kn_j, \text{ for } k \in \mathbb{N}, \text{ then}$$

$$\text{tr}(T_j^m) = \left(2^k - \left[\begin{array}{c} k \\ \frac{k}{2} \end{array} \right] \right) + \left[\begin{array}{c} m \\ \frac{m}{2} \end{array} \right],$$

$$(3) \text{ if } m = k_1 n_j + k_2, \text{ for } k_1 \in \mathbb{N} \text{ and } 1 \leq k_2 < n_j, \text{ then}$$

$$\text{tr}(T_j^m) = \left(2^{k_1} - \left[\begin{array}{c} k_1 \\ \frac{k_1}{2} \end{array} \right] \right) + \left[\begin{array}{c} m \\ \frac{m}{2} \end{array} \right].$$

□

We will finish this chapter with the following remark ;

Remark 2.1. *Let $G = \langle X : R \rangle$ be a finitely presented group with its generator set $X = \{x_1, \dots, x_N\}$ and the relation $R = \{r_1, \dots, r_M\}$ and let's fix a generator $x_j \in X$ and the corresponding block operator $T_j = x_j + x_j^{-1}$. Suppose there exist $r_t \in R$ and $n_j \in \mathbb{N}$ such that $r_t = x_j^{n_j}$. Then, by (2) and (3) of the previous theorem,*

$$\text{tr}(T_j^m) = \begin{cases} \left[\begin{array}{c} m \\ \frac{m}{2} \end{array} \right] & \text{if } m < n_j \\ \left(2^{k_1} - \left[\begin{array}{c} k_1 \\ \frac{k_1}{2} \end{array} \right] \right) + \left[\begin{array}{c} m \\ \frac{m}{2} \end{array} \right] & \text{if } m \geq n_j, \end{cases}$$

where $m = k_1 n_j + k_2$, for $k_1 \in \mathbb{N}$ and $k_2 \in \mathbb{N} \cup \{0\}$. $\square$

2.2. Identically Distributedness.

By the previous section, we have that if $G = \langle X : R \rangle$ is a finitely presented group and if $x \in X$, then

- (i) if there exists $n \in \mathbb{N}$ such that $x^n = r$, for $r \in R$, then

$$tr(T^m) = \begin{cases} \left\lfloor \frac{m}{2} \right\rfloor & \text{if } m < n_j \\ \left(2^{k_1} - \left\lfloor \frac{k_1}{2} \right\rfloor \right) + \left\lfloor \frac{m}{2} \right\rfloor & \text{if } m \geq n_j, \end{cases}$$

where $m = k_1 n + k_2$, for $k_1 \in \mathbb{N}$ and $k_2 \in \mathbb{N} \cup \{0\}$, and

- (ii) if there is no $n \in \mathbb{N}$ such that $r = x^n$, for all $r \in R$, then

$$tr(T^m) = \left\lfloor \frac{m}{2} \right\rfloor, \text{ for all } m \in \mathbb{N}.$$

The above formulæ directly proves the following theorem ;

Theorem 2.5. *Let $G_i = \langle X_i : R_i \rangle$ be finitely presented groups, for $i = 1, 2$, and assume that $x_i \in X_i$ are generators of G_i , for $i = 1, 2$. If there are relators $r_i \in R_i$ and $n \in \mathbb{N}$ such that $r_i = x_i^n$, for all $i = 1, 2$, then the block operators $(x_i + x_i^{-1})$ of the group von Neumann algebras $L(G_i)$, $i = 1, 2$, are identically distributed.*

Proof. By (i) and (ii) in the previous paragraph, case by case, we can get that

$$tr(T_1^m) = tr(T_2^m), \text{ for all } m \in \mathbb{N}.$$

■

In the above theorem, G_1 and G_2 are not necessarily distinct. i.e., suppose that we have a finitely presented group $G = \langle X : R \rangle$ and its generators x_1, x_2 in X satisfying that x_1^n and x_2^n are relators in R . Then the block operators $(x_1 + x_1^{-1})$ and $(x_2 + x_2^{-1})$ are identically distributed in the group von Neumann algebra, $L(G)$.

The following theorem is proved, similarly ;

Theorem 2.6. *Let $G_i = \langle X_i : R_i \rangle$ be finitely presented groups, for $i = 1, 2$, and assume that $x_i \in X_i$ are generators of G_i , for $i = 1, 2$. Suppose that there is no numbers $n_1, n_2 \in \mathbb{N}$ such that $x_i^{n_i} \in R_i$, for $i = 1, 2$, then the block operators $x_i + x_i^{-1}$ in $L(G_i)$ are identically distributed. $\square$*

Reference

- [1] A. Nica, R-transform in Free Probability, IHP course note, available at www.math.uwaterloo.ca/~anica.
- [2] A. Nica and R. Speicher, R-diagonal Pair-A Common Approach to Haar Unitaries and Circular Elements, (1995), www.mast.queensu.ca/~speicher.
- [3] B. Solel, You can see the arrows in a Quiver Operator Algebras, (2000), preprint.
- [4] D. Shlyakhtenko, Some Applications of Freeness with Amalgamation, J. Reine Angew. Math, 500 (1998), 191-212.
- [5] D. Voiculescu, K. Dykemma and A. Nica, Free Random Variables, CRM Monograph Series Vol 1 (1992).
- [6] D. Voiculescu, Operations on Certain Non-commuting Operator-Valued Random Variables, Astérisque, 232 (1995), 243-275.
- [7] D. Shlyakhtenko, A-Valued Semicircular Systems, J. of Funct Anal, 166 (1999), 1-47.
- [8] D.W. Kribs and M.T. Jury, Ideal Structure in Free Semigroupoid Algebras from Directed Graphs, preprint.
- [9] D.W. Kribs and S.C. Power, Free Semigroupoid Algebras, preprint.
- [10] I. Cho, Graph W^* -Probability Theory, (2004), Preprint.
- [11] I. Cho, Random Variables in Graph W^* -Probability Spaces, (2004), Preprint.
- [12] I. Cho, Amalgamated Semicircular Systems in Graph W^* -Probability Spaces, (2004), Preprint.
- [13] I. Cho, Weighted Graph W^* -Probability Spaces, (2004), Preprint.
- [14] I. Cho, The Moment Series of the Generating Operator of $L(F_2)*_{L(F_1)}L(F_2)$, (2003), Preprint.
- [15] I. Grossman and W. Magnus, Groups and Their Graphs, MAA (1964), ISBN-0-88385-600-X.
- [16] K. J. Horadam, The Word Problem and Related Results for Graph Product Groups, Proc. AMS, vol. 82, No 2, (1981) 157-164.
- [17] J. Lauri and R. Scapellato, Topics in Graph Automorphisms and Reconstruction, London Math Soc. Student No. 54, (1995), Cambridge Univ. Press.

[18] P. Biane and R. Speicher, Stochastic Calculus with respect to Free Brownian Motion and Analysis on Wigner Space, Prob. Theory Relat. Fields 112, (1998) 378-409.

[19] R. Speicher, Combinatorial Theory of the Free Product with Amalgamation and Operator-Valued Free Probability Theory, AMS Mem, Vol 132 , Num 627 , (1998).

[20] R. Speicher, Combinatorics of Free Probability Theory IHP course note, available at www.mast.queensu.ca/~speicher.

[21] S. Hermiller and J. Meier, Algorithms and Geometry for Graph Products of Groups, J of Algebra 171, (1995) 230-257.

(A. One and A. Two) AUTHOR ONE TWO ADDRESS LINE 1, AUTHOR ONE TWO ADDRESS LINE 2
E-mail address, A. One: `aone@aoneinst.edu`